

Fluorine Abundances in the Orion Nebula Cluster

Katia Cunha

Observatório Nacional, Rua General José Cristino 77, 20921-400, São Cristóvão, Rio de Janeiro, Brazil; katia@on.br

Verne V. Smith

National Optical Astronomy Observatory, P.O. Box 26732, Tucson, AZ 85726, USA; vsmith@noao.edu

ABSTRACT

This is a pilot study using cool dwarfs as sources with which to probe fluorine abundances via HF. This molecule is detected for the first time in young K-M dwarf members of an OB association. The targets are three low-mass stars (JW22, JW163 and JW433) belonging to the Orion Nebula Cluster. The target stellar parameters were derived to be $T_{\text{eff}}=3650\text{K}$, 4250K , and 4400K , and $\log g = 3.5$, 3.4 and 3.6 , with corresponding stellar masses of 0.4 , 0.6 and $0.7M_{\odot}$, for JW22, JW163 and JW433, respectively. Fluorine, oxygen, and carbon abundances were derived from the HF(1–0) R9 line along with samples of OH and CO vibration-rotation lines present in high-resolution infrared spectra observed with the Phoenix spectrograph on the Gemini South Telescope. The fluorine and oxygen results obtained for these targets, still in the pre-main-sequence stage of evolution, agree well with the general trend defined for the Milky Way disk; the latter being deduced from observations of more evolved giant stars. In addition, the carbon and oxygen abundances obtained for the studied stars overlap results from previous studies of the more massive OB stars and FG dwarf members of the Orion Nebula Cluster. We conclude from this agreement that the fluorine abundances derived for the Orion K-M dwarfs (when there is no conspicuous evidence of disks) can be considered a good representation of the current fluorine abundance value for the Milky Way disk.

Subject headings: nucleosynthesis—stars: abundances

1. Introduction

The chemical elements are produced via a variety of nuclear and high-energy processes associated with stars of various masses, cosmic rays, and the big bang itself. One inter-

esting process, whose importance within the overall scheme of nucleosynthesis is not yet well-quantified, involves the inelastic scattering of neutrinos off of nuclei and is referred to as neutrino nucleosynthesis, or the ν -process. Neutrino nucleosynthesis was discussed by Domogatsky, Eramzhyan, & Nadyozhin (1978), who focussed mainly on its role in the production of the lightest elements, such as ^2H , ^3He , ^7Li , or ^{11}B . Later, Woosley & Haxton (1988) expanded on the role of the ν -process in the production of fluorine and subsequently, Woosley et al. (1990) published predicted yields for a number of species, including Li and F. The neutrinos involved in the ν -process are produced as a result of the gravitational collapse of a stellar core to a neutron star during a supernova of Type II (SN II). The ν -process may result in significant synthesis of certain low-abundance isotopes that lie one mass unit below abundant nuclei, such as ^{12}C giving rise to ^{11}B , or ^{20}Ne resulting in ^{19}F via neutrino-induced spallation. It is the association of ^{19}F , the only stable isotope of fluorine, with the ν -process that makes its abundance of interest as a potential probe of neutrino nucleosynthesis.

In addition to the ν -process, ^{19}F is possibly produced during He-burning thermal pulses on the asymptotic giant branch (AGB), as first suggested by Forestini et al. (1992). Meynet & Arnould (2000) also pointed out that Wolf-Rayet stars, via the same sets of reactions as in the AGB stars, may also produce significant amounts of ^{19}F , with the caveat that large dM/dt stellar winds must remove a significant amount of mass, exposing the inner layers where fluorine is produced before it is burned away. Observations of fluorine abundances, spanning a range of stellar metallicities and across a variety of stellar populations, are needed in order to sort out relative contributions from AGB stars, Wolf-Rayet stars, or neutrino nucleosynthesis.

In this paper, we use high-resolution IR spectra from Phoenix on Gemini-South to derive ^{19}F abundances (from HF), along with ^{12}C abundances (from CO), and ^{16}O abundances (from OH) in three low-mass members of the Orion Nebula cluster. Carbon and oxygen provide comparison abundances to those derived from fluorine. Both C and O have been studied previously in Orion members by Cunha & Lambert (1994); their study analyzed hot O and B stars. In addition, some oxygen abundances in Orion-member F and G main-sequence and pre-main-sequence stars have been published by Cunha, Smith, & Lambert (1998). These previous studies provide a background on which the abundances derived from K and M type stars can be compared. Apart from the interest in the carbon and oxygen abundances in young stellar systems, C and O provide important comparisons for the fluorine abundances. The Orion Nebula cluster stars are very young and will reflect the current Galactic F/O ratio in the disk. Also, previous stellar ^{19}F abundances (derived from HF) have come only from red giants (Jorissen, Smith, & Lambert 1992; Cunha et al. 2003), while here we undergo a pilot study to use cool dwarf stars as sources with which to probe fluorine abundances.

2. Observations

The target stars for this study are K and M dwarfs selected from various studies of the Orion Nebula cluster by Hillenbrand (1997), Hillenbrand & Hartmann (1998), Hillenbrand et al. (1998), and Rhode, Herbst, & Mathieu (2001). In particular, the stars were selected for high membership probability ($P \geq 93\%$), to be cool enough for the HF molecular line to be detectable (cooler than T_{eff} of 4500K), and to have relatively low values of projected rotational velocities ($v \sin(i) \leq 15 \text{ km s}^{-1}$). The selected targets were such that two of them (JW22 and JW433) do not show evidence of flux from a circumstellar disk, while JW163 shows such evidence (primarily from near-IR color excesses from Hillenbrand et al. 1998). Our sample, although small, will allow us to ascertain possible effects of disk emission on an abundance analysis of the stellar photosphere in these young stars.

High-resolution infrared spectra of these low-mass stars were obtained with the Gemini South Telescope plus the Phoenix spectrograph (Hinkle et al. 1998) during observing runs in classical mode on 27-28 December 2002 and 20 December 2004. The observed data are single order echelle spectra with a resolution $R = \lambda / \Delta\lambda = 50,000$, corresponding to a resolution element of ~ 4 -pixels. In order to obtain two separate spectral regions, one around the H-band (at $\sim 15,500\text{\AA}$, containing ^{16}OH lines) and another around the K-band (at $\sim 23,400\text{\AA}$, containing the molecular HF(1–0) R9 line and numerous $^{12}\text{C}^{16}\text{O}$ lines), two grating tilts were needed. JW22 was observed only in the K-band region.

The Orion targets are relatively bright in the near-infrared so that the exposure times needed to obtain co-added spectra with signal-to-noise ratios in excess of ~ 50 – 100 were not exceedingly long: for each grating tilt, four target exposures of 15 minutes each were obtained. We adopted the strategy of varying the target position on the slit (positions a-b-b-a) and subtracting one image from another in order to remove dark current and fixed-pattern noise. In addition, for those observations corresponding to the spectral region of the K-band, we also observed hot stars at roughly the same air masses as the target stars. These hot star spectra were needed in order to remove telluric lines from the target star spectra. All the obtained data were reduced to one-dimensional spectra using standard IRAF routines. More details on the observations and reduction procedures adopted can be found in Smith et al. (2004).

Phoenix spectra are shown in Figure 1 for two of the target stars, with the two spectra shifted vertically in relative flux. The spectra plotted in this figure were obtained in the december 2002 Gemini run. The top panel shows the spectra in the H band, while the bottom spectra are in the K band. Some of the prominent absorption lines are identified in the figure, with the $15550\text{\AA}$ region showing a number of OH lines and the $23400\text{\AA}$ spectrum containing numerous $^{12}\text{C}^{16}\text{O}$ lines.

3. Stellar Parameters

The effective temperatures for the studied stars were derived from their photometric colors (V-I) and (V-K) and by adopting the empirical calibrations from Bessel, Castelli & Plez (1998). (V-I) and (V-K) colors are considered to be good temperature indicators for cool stars and have little sensitivity to gravity for the effective temperature range considered in this study. The adopted photometric magnitudes are presented in Table 1. We obtained V-magnitudes and (V-I) colors for our targets from the work by Hillenbrand (1997) that studied the stellar population comprising the Orion Nebula cluster. The target star K-magnitudes were obtained from the Two Micron All Sky Survey (2MASS) database and corrected to the Bessel et al. system by means of the transformation equations presented in Carpenter (2001). De-reddened colors were computed by adopting $A_v=0$ for JW163, $A_v=1.01$ mag for JW433, and $A_v=0.8$ for JW22 (all values are taken from Hillenbrand 1997).

Surface gravities ($\log g$'s) for the target stars were calculated from the following standard relation:

$$g/g_{\odot} = (M/M_{\odot}) (T_{eff}/T_{\odot})^4 (L/L_{\odot})^{-1}.$$

The stellar luminosities were taken from Hillenbrand (1997; $\log L/L_{\odot} = 0.22, 0.43$, and -0.34 for JW163, JW433, and JW22, respectively). These were derived assuming a distance to the Orion Nebula of 470 pc. As discussed in Hillenbrand (1997), the adopted stellar luminosities are estimated to have an uncertainty of 0.12 in $\log L/L_{\odot}$. The effective temperatures adopted to derive $\log g$'s were those listed in Table 1. Stellar masses could then be estimated from placement of the targets on an HR diagram and noting their position relative to evolutionary tracks computed by D'Antona & Mazzitelli (1994). Surface gravities were then computed for the sample stars. The derived masses and $\log g$'s are presented in Table 1. From the uncertainties in the adopted stellar luminosities and effective temperatures we estimate that our derived surface gravities have a maximum uncertainty of ~ 0.27 dex.

4. Analysis

LTE fluorine, as well as carbon and oxygen abundances were obtained from spectrum synthesis using the synthesis code MOOG (Sneden, 1973). The model atmospheres employed in the analysis were calculated with the ATLAS9 code (Kurucz 1994) for solar metallicity, adopting a mixing length $l=1.25 H_p$ and without convective overshooting.

4.1. Fluorine, Carbon and Oxygen Abundances

Fluorine abundances can be derived from the vibration-rotation lines of HF present in the spectral region around the K-band. In this study, we used the HF(1–0) R9 line at 23357.661 Å. In this same IR spectral window where HF was observed, several other molecular lines that correspond to $^{12}\text{C}^{16}\text{O}$ first-overtone vibration rotation lines (3–1 vibration series) are also found and these can be used to obtain the carbon abundances, given that the oxygen abundances are known from other transitions (such as OH). Synthetic spectra were calculated across the entire spectral region of the observed Phoenix spectra, or from ~ 23340 Å to 23430 Å. The linelist adopted was the same as in our previous study of fluorine in red-giants of the Large Magellanic Cloud (see Cunha et al. 2003). The CO lines are typically used to derive the microturbulence in cool giants, where they are much stronger (such as in Smith et al. 2002). Although in the parameter range studied here, the CO lines become weaker, they are still sensitive to the microturbulence and were used in order to estimate this parameter. The adopted values for the stars vary between $\xi=0.5$ and 1.0 km/s (Table 1).

Figure 2 shows the synthetic and observed spectra for all three target stars near the HF line. Three synthetic spectra, each with a different fluorine abundance, are plotted in each panel in order to illustrate the sensitivity of the HF line to the fluorine abundance. Just to the blue of the HF(1–0) R9 line is a stellar photospheric line that is probably due to H_2O . This same unidentified line is clearly visible (in the same approximate relative strength) in the solar sunspot spectrum published by Wallace & Livingston (1992). They identify a number of other H_2O lines in this spectral region that have similar strength. Water is not included in the synthesis linelist here, and the offending line blends just a small fraction of the blue edge of the HF line and does not affect significantly the derived ^{19}F abundance. If triatomic H_2O is responsible for this line, then it should be much weaker in giant stars of similar effective temperature, and this can be seen in the spectral atlas of Arcturus (Hinkle, Wallace, & Livingston 1995); although having a similar effective temperature to JW163 or JW433, this feature is not detectable in Arcturus, even though the HF line is clearly visible. Indeed, a tentative identification has been provided by Hinkle (2004) as H_2O (021)[14,5,9]–(010)[13,11,2] at an air wavelength of about 23356.4 Å.

Oxygen abundances can be measured from first-overtone vibration-rotation OH lines present in the H-band region. Three isolated OH lines at 15560, 15569 and 15572 Å were synthesized to derive oxygen abundances. The studied OH lines and adopted linelist were the same as in Smith et al. (2002). These lines are expected to be reliable oxygen abundance indicators and relatively free from non-LTE effects but to suffer modestly (~ 0.1 dex) from 3d-effects in solar-type stars (Asplund 2005). However, at the relatively low effective temperatures of the stars studied here, the oxygen abundance differences caused by the adoption

of one-dimensional model atmospheres, instead of three-dimensional model atmospheres, are expected to be smaller than at higher effective temperatures (around 6000K; Nissen et al. 2002).

Overall, the derived abundances in JW163, relative to JW433 and JW22, are systematically lower for the studied elements, with ^{12}C 0.30-0.40 dex lower, ^{16}O 0.25 dex lower, and ^{19}F 0.10 dex lower. Although it is possible that this difference reflects an underlying abundance spread in the Orion Nebula cluster, it is more likely that the presence of a more substantial circumstellar disk in JW163 results in the photospheric absorption lines being filled in by disk emission. It is worth noting that in the (V-I)-(V-K) plane as shown in Figure 11 of Bessell et al. (1998), both JW433 and JW22 fall right on the sequence defined by normal dwarf stars, while JW163 is displaced several tenths of a magnitude towards redder colors from the standard dwarf relation. In JW433 and JW22, with little if any measurable disk presence, the abundances are near-solar: $A(^{12}\text{C})=8.30$, $A(^{16}\text{O})=8.80$, and $A(^{19}\text{F})=4.65$ for JW433, while for JW22 the abundances are $A(^{12}\text{C})=8.40$ and $A(^{19}\text{F})=4.61$. Moreover, as will be discussed in Section 5, the carbon (for JW22 and JW433) and oxygen (for JW433) abundances overlap nicely with those values derived from the much hotter B-star members of Orion Id (the sub-association to which the Orion Nebula cluster belongs) by Cunha & Lambert (1994). The results from JW163, along with JW22 and JW433 illustrate the usefulness of using these cool dwarfs for abundance studies, even those still not on the main sequence, provided that targets are selected carefully, taking into account their circumstellar environments.

4.2. Abundance Uncertainties

The final abundances derived for carbon, oxygen and fluorine are presented in Table 1. The expected uncertainties in the derived effective temperatures and surface gravities are 100K and 0.3 dex, respectively. The errors in the microturbulence parameters are estimated to be around 0.5 km/s. The sensitivity of the derived C, O and F abundances to changes in temperature, gravity and microturbulence for two target stars (the coolest and the hottest in our sample) are presented in Table 2; the total errors (ΔA) in the derived abundances in each case are listed in column 6. Overall, the studied transitions of HF, CO and OH are weak and relatively insensitive to the microturbulence; the exception being the CO lines that are moderately strong at T_{eff} 's around 3650K. The fluorine abundances are mostly sensitive to effective temperature, but also, to a lesser extent, to the surface gravity. Concerning OH, the abundance errors are quite insensitive to changes in surface gravity and microturbulence. The total abundance errors for oxygen are dominated by T_{eff} uncertainties. (OH abundances were

not calculated for JW22 in this study.) The uncertainties in the derived carbon abundances are shown to have different sensitivities at the "cool" and "hot" regimes studied here. These are more uncertain around 3600K when compared to 4500K.

5. Discussion

Our larger goal is to investigate the behavior of fluorine abundances with metallicity, which is taken here to be represented by the element oxygen, as this element is virtually a pure product of nucleosynthesis in massive stars that end their evolution as core-collapse SN II. In particular, in this study we focus on the fluorine and oxygen contents of the pre-main-sequence stellar members of the Orion Nebula Cluster.

As mentioned in the introduction carbon and oxygen abundances derived from molecular transitions (CO and OH) can be used as benchmarks with which to gauge the fluorine abundances obtained from HF. A comparison of carbon and oxygen abundances derived here for JW22 and JW433 with abundances from Cunha & Lambert (1994) and Cunha et al. (1998) is in order; we ignore JW163 in this comparison as there is evidence that the absorption lines may be affected measurably by emission from a circumstellar disk. The Orion Association is divided into four subgroups, labelled Ia, Ib, Ic, and Id. The subgroups are distributed roughly by age, with Ia being the oldest, Id the youngest, and an age difference of about 10Myr. The Orion Nebula stars analyzed in this study are members of Id. The abundances in the two stars are $A(C)=8.40$ for JW22, and $A(C)=8.30$ and $A(O)=8.80$ for JW433. For comparison, Cunha & Lambert (1994) studied three OB stars in Id and their average (NLTE) abundances are $A(C)=8.25$ and $A(O)=8.85$: the agreement between two K-M dwarfs and three O-B stars is excellent. In addition, Cunha et al. (1998) have one F-dwarf that is a Id member and its oxygen abundance is $A(O)=8.86$: again, excellent agreement. Oxygen and carbon abundances in the OB stars come from analyses of O II and C II lines, while oxygen in the F dwarf comes from an analysis of O I lines. The K-M dwarfs use molecular OH and CO lines. This abundance comparison across a large range in T_{eff} utilizing a variety of spectroscopic signatures provides evidence that the abundance analysis of the low-mass stars studied here is robust. We proceed to a discussion of the fluorine abundances.

Historically, ^{19}F abundances were derived initially in a sample of galactic K, M, S, and C red-giants, with metallicities mostly near solar, by Jorissen et al. (1992); this study was restricted to fairly bright stars in the IR. More recently, access to high spectral resolution in the IR using the Phoenix spectrograph on the Gemini-South 8.1m telescope has allowed fluorine abundances to be measured in fainter samples of cool stars, such as in the Large

Magellanic Cloud (LMC) or in the globular cluster ω Centauri (Cunha et al. 2003). Renda et al. (2004) used the abovementioned ^{19}F abundances as comparisons to chemical evolution models. They conclude that both Wolf-Rayet and AGB stars are significant sources of ^{19}F , with the Wolf-Rayet stars dominating at high metallicities. At low metallicities (less than roughly -0.8 dex) synthesis via the ν process in SN II dominates fluorine production. Renda et al. (2004) were the first to show that all three ^{19}F sources can be significant contributors, but with differing relative contributions depending on the metallicity and star formation history of the stellar population. In addition to abundances derived from stellar HF lines, Federman et al. (2005) have obtained detections of interstellar F I lines (at 955Å) along two lines-of-sight into the Cep OB2 association using the Far Ultraviolet Spectroscopic Explorer (FUSE). Their derived column densities of F and O indicate a near-solar F/O ratio, with some small depletion effects.

The Orion stars analyzed here are of particular interest as they represent a different type of star with which to probe fluorine when compared to previous studies since Jorissen et al. (1992) sampled only evolved stars. Our work adds three stars of higher surface gravity to the sample of near-solar metallicity field stars and their ^{19}F and ^{16}O abundances are shown in Figure 3 in the form of [F/O] versus A(O). The point represented by JW163 should be viewed with caution as we have noted already that its abundance results have probably been influenced by the presence of a circumstellar disk. Although both O and F could have been affected about the same with the result that the ratio of F/O is changed relatively less. Also, we adopt the same oxygen abundance for JW22 as we derived for JW433 because the 15550Å region with the OH lines was not observed for JW22; however its carbon abundance is the same, within the errors, as found for JW433 and there is no evidence that the Orion stars exhibit large variations in their total C/O values. The Jorissen et al. (1992) K and M giants are also plotted (with the abundances in these stars re-derived by Cunha et al. (2003)); the solar values in Figure 3 are taken to be A(O)= 8.69 and A(F)= 4.55. Note that the Orion dwarf abundances overlap very nicely with the Jorissen et al. giants.

The stellar abundances that are plotted in Figure 3 represent all the available fluorine abundances that have been measured for members of the Milky Way so far (excluding self-polluted AGB stars). Abundances for LMC red giants from Cunha et al. (2003) are not shown here, as the star formation history and corresponding chemical evolution in the LMC may be quite different from the Milky Way. The solid curve in Figure 3 is a chemical evolution model from Renda et al. (2004) for the Milky Way. This particular curve represents their MWc model that includes ^{19}F yields from all three stellar production sites: SN II with the ν -process, Wolf-Rayet stars, and AGB stars. The Orion Nebula cluster F and O abundances help anchor the Milky Way values at high metallicity, which are fit nicely by the Renda et al. (2004) model with all three sources. The low values of [F/O] in the ω Cen stars were

discussed by both Cunha et al. (2003) and Renda et al. (2004) in the context of the unusual star formation history and chemical evolution within this self-enriched stellar system that has been captured by the Galaxy.

6. Conclusions

We detect the HF molecular line at 23400Å in K-M dwarfs and present the first fluorine abundance measurements in young stars still in the pre-main sequence phase of evolution. These targets are members of the Orion association, in particular from the Orion Nebula cluster, with masses around 0.4-0.8 $M_{\odot}$. Our results for JW433 and JW22, which do not show substantial emission from a circumstellar disk, reveal abundances of C and O that are in excellent agreement with those that are found in B-type stellar members of Orion Id (Cunha & Lambert 1994). This agreement bolsters confidence in the quantitative spectroscopic abundance analyses techniques used in the hotter B stars, as well as the cool K and M dwarfs. Moreover, JW433 and JW22 also display abundance patterns in agreement with the general behavior of fluorine versus oxygen for the Milky Way disk that has been established from stars in a distinctly different evolutionary state. All previous F abundances had been derived from studies of red giants. The overlap in these abundances provides an important confirmation for the relation of the fluorine abundance with oxygen in the Disk.

In the future, larger samples of K and M dwarfs, as well as L and T dwarfs, can be used to measure fluorine and oxygen abundances both in the field and in the nearer star-forming regions.

Based on observations obtained at the Gemini Observatory, which is operated by the Association of Universities for Research in Astronomy, Inc., under a cooperative agreement with the NSF on behalf of the Gemini partnership: the National Science Foundation (United States), the Particle Physics and Astronomy Research Council (United Kingdom), the National Research Council (Canada), CONICYT (Chile), the Australian Research Council (Australia), CNPq (Brazil), and CONICRT (Argentina), as program GS-2002B-Q-2. This work is also supported in part by the National Science Foundation through AST03-07534 (VVS) and NASA through NAG5-9213 (VVS).

Allende Prieto, C., Lambert, D. L., & Asplund, M. 2000, *ApJ*, 573, L137

Asplund, M. 2005, to appear in *Cosmic Abundances as Records of Stellar Evolution and Nucleosynthesis*, ASP Conference Series, eds. F. N. Bash & T. G. Barnes (San Francisco:ASP)

Bessel, M.S., Castelli, F. & Plez, B. 1998, *A&A*, 333, 231

Carpenter, J. M. 2001, *ApJ*, 121, 2851

Cunha, K., Smith, V.V., Lambert, D.L. & Hinkle, K.H. 2003, *AJ*, 126, 1305

D’Antona, F. & Mazzitelli, I., 1994, *ApJS*, 90, 467

Domogatskii, G. V., Eramzhian, R. A., & Nadezhin, D. K. 1978, *Ap.& Space Sci.*, 58, 273

Federman, S. R., Sheffer, Y., Lambert, D. L., & Smith, V. V. 2005, *ApJ*, in press (astro-ph/0410362)

Forestini, M., Goriely, S., Jorissen, A., & Arnould, M. 1992, *A&A*, 261, 157

Hillenbrand, L. A. 1997, *AJ*, 1733, 1768

Hillenbrand, L. A., & Hartmann, L. W. 1998, *ApJ*, 492, 540

Hillenbrand, L. A., Strom, S. E., Calvet, N., Merrill, K.M., Gatley, I., Makidon, R.B., Meyer, M.R., & Skrutskie, M.F., 1998, *AJ*, 116, 1816

Hinkle, K. H. 2004, private communication

Hinkle, K. H., Wallace, L., & Livingston, W. 1995, *Infrared Atlas of the Arcturus Spectrum, 0.9–5.3 μ m* (San Francisco:ASP)

Hinkle, K. H., Cuberly, R., Gaughan, N., Heynssens, J., Joyce, R., Ridgway, S., Schmitt, P., & Simmons, J. E. 1998, *Proc. SPIE*, 3354, 810

Jorissen, A., Smith, V.V., & Lambert, D.L., *A&A*, 261, 164

Kurucz, R. L. 1994, CD-ROM 19 (Cambridge:SAO)

Meynet, G., & Arnould, M. 2000, *A&A*, 355, 176

Nissen, P.E., Primas, F., Asplund, M., & Lambert, D.L. 2002, *A&A* 390, 235

Renda, A. et al. 2004 , *MNRAS*, 354, 575

Rhode, K.L., Herbst, W., & Mathieu, R.D. 2001, *AJ*, 122, 3258

Snedden, C. 1973, *ApJ*, 184, 839

Smith, V.V., Hinkle, K.H., Cunha, K., Plez, B., Lambert, D.L., Pilachowski, C.A., Barbuy, B., Melendez, J., Balachandran, S., Bessel, M.S., Geisler, D.P., Hesser, J.E., & Winge, C., 2002, *AJ*, 124, 3241

Wallace, L., & Livingston, W. 1992, *An Atlas of a Dark Sunspot Umbral Spectrum from 1970 to 8640 cm⁻¹ (1.16 to 5.1 μ m)*, N.S.O. Technical Report #92-001

Woosley, S.E., Haxton, W.C., 1988, *Nature*, 334, 45

Woosley, S.E., Hartmann, D.H., Hoffman, R.D., Haxton, W.C., 1990, *ApJ*, 356, 272

Table 1. Stellar Parameters and Abundances

Star	V_0	I_0	K_0	T_{eff}	$\text{Log } g$	ξ	$M/M_{\odot}$	$A(^{12}\text{C})$	$A(^{16}\text{O})$	$A(^{19}\text{F})$
JW 22	15.5	13.4	11.3	3650	3.6	1.0	0.4	8.40	—	4.61
JW 163	13.1	11.8	10.2	4250	3.5	0.5	0.6	8.00	8.55	4.53
JW 433	12.5	11.3	9.6	4400	3.4	0.8	0.7	8.30	8.80	4.65

Note. — $A(X) = \log[n(X)/n(\text{H})] + 12$.

Table 2. Abundance Uncertainties

Star	Element	$\Delta T_{\text{eff}}(+100\text{K})$	$\Delta \text{Log } g(+0.3 \text{ dex})$	$\Delta \xi(+0.5 \text{ km/s})$	ΔA
JW 22	ΔC	0.08	0.0	0.10	0.13
JW 22	ΔF	0.08	0.05	0.0	0.09
JW 433	ΔC	0.15	0.15	0.0	0.21
JW 433	ΔO	0.10	0.02	0.0	0.10
JW 433	ΔF	0.13	0.05	0.0	0.14

Fig. 1.— Observed Phoenix spectra for the target stars JW163 and JW433 centered at 15545 Å (top panel) and 23390 Å. The spectra have been separated vertically, while prominent absorption lines are indicated.

Fig. 2.— Observed and synthetic spectra for the three target stars in the HF region. The best-fit fluorine abundances (represented by solid lines) are those presented in Table 1. The synthetic spectra represented by the dashed lines were calculated with ± 0.2 dex changes in the ^{19}F abundance. Note the blending feature just blueward of the HF(1-0) R9 line, which is identified tentatively here as H_2O .

Fig. 3.— Values of $[\text{F}/\text{O}]$ are plotted versus oxygen abundances for field K and M giants from Cunha et al. (2003) and the three Orion Nebular cluster stars. The solar symbol is shown and the solid line illustrates the predictions from one chemical evolution model by Renda et al. (2004) that includes all three suggested sources for ^{19}F production. Note that the Orion Nebula stars fall within the scatter defined by the red giants of the Milky Way field.





